

Protecting Water Quality *from* **AGRICULTURAL RUNOFF**

Clean Water Is Everybody's Business

The United States has more than 330 million acres of agricultural land that produce an abundant supply of food and other products. American agriculture is noted worldwide for its high productivity, quality, and efficiency in delivering goods to the consumer. When improperly managed however, activities from working farms and ranches can affect water quality.

In the 2000 *National Water Quality Inventory*, states reported that agricultural nonpoint source (NPS) pollution is the leading source of water quality impacts on surveyed rivers and lakes, the second largest source of impairments to wetlands, and a major contributor to contamination of surveyed estuaries and ground water. Agricultural activities that cause NPS pollution include poorly located or managed animal feeding operations; overgrazing; plowing too often or at the wrong time; and improper, excessive, or poorly timed application of pesticides, irrigation water, and fertilizer.

Pollutants that result from farming and ranching include sediment, nutrients, pathogens, pesticides, metals, and salts. Impacts from agricultural activities on surface water and ground water can be minimized by using management practices that are adapted to local conditions. Many practices designed

What Is Nonpoint Source Pollution?

Nonpoint source (NPS) pollution, unlike pollution from point sources such as industrial and sewage treatment plants, comes from many diffuse sources. Polluted runoff is caused by rainfall or snowmelt moving over and through the ground. As the runoff moves, it picks up and carries away natural and human-made pollutants, finally depositing them into watersheds through lakes, rivers, wetlands, coastal waters, and even our underground sources of drinking water.

Did you know that runoff from farms is the leading source of impairments to surveyed rivers and lakes?

to reduce pollution also increase productivity and save farmers and ranchers money in the long run.

There are many government programs available to help farmers and ranchers design and pay for management approaches to prevent and control NPS pollution. For example, over 40 percent of section 319 Clean Water Act grants have been used to control NPS pollution from working farms and ranches. Also, many programs funded by the U.S. Department of Agriculture and by states provide cost-share, technical assistance, and economic incentives to implement NPS pollution management practices. Many local organizations and individuals have come together to help create regional support networks to adopt technologies and practices to eliminate or reduce water quality impacts caused by agricultural activities.

Sedimentation

The most prevalent source of agricultural water pollution is soil that is washed off fields. Rain water carries soil particles (sediment) and dumps them into nearby lakes or streams. Too much sediment can cloud the water, reducing the amount of sunlight that reaches aquatic plants. It can also clog the gills of fish or smother fish larvae.

In addition, other pollutants like fertilizers, pesticides, and heavy metals are often attached to the soil particles and wash into the water bodies, causing algal blooms and depleted oxygen, which is deadly to most aquatic life. Farmers and ranchers can reduce erosion and sedimentation by 20 to 90 percent by applying management practices that control the volume and flow rate of runoff water, keep the soil in place, and reduce soil transport.

Nutrients

Farmers apply nutrients such as phosphorus, nitrogen, and potassium in the form of chemical fertilizers, manure, and sludge. They may also grow legumes and leave crop residues to enhance production. When these sources exceed plant needs, or are applied just before it rains, nutrients can wash into aquatic ecosystems. There they can cause algae blooms, which can ruin swimming and boating opportunities, create foul taste and odor in drinking water, and kill fish by removing oxygen from the water. High concentrations of nitrate in drinking water can cause methemoglobinemia, a potentially fatal disease in infants, also known as blue baby syndrome. To combat nutrient losses, farmers can implement nutrient management plans that help maintain high yields and save money on fertilizers.



Animal Feeding Operations

By confining animals in small areas or lots, farmers and ranchers can efficiently feed and maintain livestock. But these confined areas become major sources of animal waste. An estimated 238,000 working farms and ranches in the United States are considered animal feeding operations, generating about 500 million tons of manure each year. Runoff from poorly managed facilities can carry pathogens such as bacteria and viruses, nutrients, and oxygen-demanding organics and solids that contaminate shellfishing areas and cause other water quality problems. Ground water can also be contaminated by waste seepage. Farmers and ranchers can limit discharges by storing and managing facility wastewater and runoff with appropriate waste management systems.


Livestock Grazing

Overgrazing exposes soils, increases erosion, encourages invasion by undesirable plants, destroys fish habitat, and may destroy streambanks and floodplain vegetation necessary for habitat and water quality filtration. To reduce the impacts of grazing on water quality, farmers and ranchers can adjust grazing intensity, keep livestock out of sensitive areas, provide

alternative sources of water and shade, and promote revegetation of ranges, pastures, and riparian zones.

Irrigation

Irrigation water is applied to supplement natural precipitation or to protect crops against freezing or wilting. Inefficient irrigation can cause water quality problems. In arid areas, for example, where rainwater does not carry minerals deep into the soil, evaporation of irrigation water can concentrate salts. Excessive irrigation can affect water quality by causing erosion, transporting nutrients, pesticides, and heavy metals, or decreasing the amount of water that flows naturally in streams and rivers. It can also cause a buildup of selenium, a toxic metal that can harm waterfowl reproduction. Farmers can reduce NPS pollution from irrigation by improving water use efficiency. They can measure actual crop needs and apply only the amount of water required. Farmers may also choose to convert irrigation systems to higher efficiency equipment.



Pesticides

Insecticides, herbicides, and fungicides are used to kill agricultural pests. These chemicals can enter and contaminate water through direct application, runoff, and atmospheric deposition. They can poison fish and wildlife, contaminate food sources, and destroy the habitat that animals use for protective cover. To reduce contamination from pesticides, farmers should use Integrated Pest Management (IPM) techniques based on the specific soils, climate, pest history, and crop conditions for a particular field. IPM encourages natural barriers and limits pesticide use and manages necessary applications to minimize pesticide movement from the field.

Farm Bill Conservation Funding

In May 2002 President Bush signed the Farm Bill, providing up to \$13 billion for conservation programs for six years. This Farm Bill represents an 80 percent increase above current levels of funding available for conservation programs designed to prevent polluted runoff. For more information, visit www.usda.gov/farmbill.

Related Publications and Web Sites

National Management Measures to Control Nonpoint Source Pollution from Agriculture

epa.gov/nps/agmm

This technical guidance and reference document is for use by state, local, and tribal managers in the implementation of nonpoint source pollution management programs. It contains information on effective, readily available, and economically achievable means of reducing pollution of surface and ground water from agriculture.

Agricultural Nonpoint Source Pollution Management Web Site

epa.gov/nps/agriculture.html

This web site features a collection of links to helpful documents, federal programs, partnerships and nongovernmental organizations that convey advice and assistance to farmers and ranchers for protecting water quality.

National Agriculture Compliance Assistance Center

epa.gov/agriculture or call toll-free: 1-888-663-2155

EPA's National Agriculture Compliance Assistance Center is the "first stop" for information about environmental requirements that affect the agricultural community.

Animal Feeding Operations (AFO) Web Sites

AFO Virtual Information Center: epa.gov/npdes/afovirtualcenter
Overview of regulations and helpful links: epa.gov/npdes/af

Funding Sources

Searchable Catalog of Federal Funding Sources for Watershed Protection

epa.gov/watershedfunding

Agricultural Management Assistance Database

www.nrcs.usda.gov/programs/ama

Clean Water Act Section 319(h) funding (epa.gov/nps/319hfunds.html) is provided to designated state and tribal agencies to implement approved nonpoint source management programs.

Environmental Quality Incentives Program (www.nrcs.usda.gov/programs/eqip) offers financial, technical, and educational assistance to install or implement structural, vegetative, and management practices designed to conserve soil and other natural resources.

Conservation Reserve and Conservation Reserve Enhancement Programs (www.fsa.usda.gov/dafp/cepd/default.htm) implemented by the U.S. Department of Agriculture provide financial incentives to encourage farmers and ranchers to voluntarily protect soil, water, and wildlife resources.

For More Information

U.S. Environmental Protection Agency
Nonpoint Source Control Branch (4503T)
1200 Pennsylvania Avenue, NW
Washington, DC 20460
epa.gov/nps